

## 10163 Storage Keepers

Randy Company has  $N$  ( $1 \leq N \leq 100$ ) storages. Company wants some men to keep them safe. Now there are  $M$  ( $1 \leq M \leq 30$ ) men asking for the job. Company will choose several from them. Randy Company employs men following these rules:

1. Each keeper has a number  $P_i$  ( $1 \leq P_i \leq 1000$ ), which stands for their ability.
2. All storages are the same as each other.
3. A storage can only be looked after by one keeper. But a keeper can look after several storages. If a keeper's ability number is  $P_i$ , and he looks after  $K$  storages, each storage that he looks after has a safe number  $U_j = P_i \div K$ . (Note:  $U_j$ ,  $P_i$  and  $K$  are all integers). The storage which is looked after by nobody will get a number 0.
4. If all the storages is at least given to a man, company will get a safe line  $L = \min U_j$
5. Every month Randy Company will give each employed keeper a wage according to his ability number. That means, if a keeper's ability number is  $P_i$ , he will get  $P_i$  dollars every month. The total money company will pay the keepers every month is  $Y$  dollars.

Now Randy Company gives you a list that contains all information about  $N$ ,  $M$ ,  $P$ , your task is give company a best choice of the keepers to make the company pay the least money under the condition that the safe line  $L$  is the highest.

### Input

The input file contains several scenarios. Each of them consists of 2 lines:

The first line consists of two numbers ( $N$  and  $M$ ), the second line consists of  $M$  numbers, meaning  $P_i$  ( $i = 1..M$ ). There is only one space between two border numbers.

The input file is ended with  $N = 0$  and  $M = 0$ .

### Output

For each scenario, print a line containing two numbers  $L(\max)$  and  $Y(\min)$ . There should be a space between them.

### Sample Input

```
2 1
7
1 2
10 9
2 5
10 8 6 4 1
5 4
1 1 1 1
0 0
```

### Sample Output

```
3 7
10 10
8 18
0 0
```