

# 10990 Another New Function

The depth of  $\phi$  value of a number is denoted by the number of steps required before it reaches 1. An example will make it very clear.

$$\begin{aligned} \phi(13) &= 12 \dots \text{step1} \\ \phi(12) &= 4 \dots \text{step2} \\ \phi(4) &= 2 \dots \text{step3} \\ \phi(2) &= 1 \dots \text{step1} \end{aligned}$$

So the depth of  $\phi(13)$  is 4. We name this function as depthphi. So we can write  $\text{depthphi}(13) = 4$ . The sum of depthphi function (SODF) takes two integers as parameter and its definition is given below:

$$\text{SODF}(m, n) = \sum_{i=m}^n \text{depthphi}(i), \quad m \leq n$$

Given the value of  $m$  and  $n$  your job is to find the value of  $\text{SODF}(m, n)$ .

**The following paragraph is extracted from Mathworld to inform you about phi function.**

The totient function  $\phi(n)$  or  $\phi(n)$ , also called Euler's totient function, is defined as the number of *positive integers*  $\leq n$  that are *relatively prime* to (i.e., do not contain any factor in common with)  $n$ , where 1 is counted as being relatively prime to all numbers. Since a number less than or equal to and *relatively prime* to a given number is called a *totative* the totient function  $\phi(n)$  can be simply defined as the number of *totatives* of  $n$ . For example, there are eight *totatives* of 24 (1, 5, 7, 11, 13, 17, 19, and 23), so  $\phi(24) = 8$ . The totient function is implemented in *Mathematica* as `EulerPhi[n]`.

## Input

The first line of the input file contains an integer  $N$  ( $0 < N < 2001$ ) which indicates how many sets of inputs are there. Each of the next  $N$  lines contains two integers  $m$  and  $n$  ( $2 \leq m \leq n \leq 2000000$ ).

## Output

For each line of input produce one line of output. This line contains an integer  $S$ , which actually denotes the value of  $\text{SODF}(m, n)$ .

## Sample Input

```
2
2 10
100000 200000
```

## Sample Output

```
22
1495105
```