

1140 Charles Frédéric Gros

Charles Frédéric Gros (CFG) has decided to disprove the Riemann hypothesis numerically. For a given integer $D > 0$ of the form $4k + 3$ and free of square prime factors, this amounts to computing the cardinality $h(D)$ of the set

$$C(D) \stackrel{\text{def}}{=} \{(a, b, c) \mid b^2 - 4ac = -D, |b| \leq a \leq c, \text{ where } b \geq 0 \text{ if } a = c \text{ or } a = |b|\}.$$

(Where a, b, c are integers.)

For instance, $C(3) = \{(1, 1, 1)\}$, $C(15) = \{(1, 1, 4), (2, 1, 2)\}$. Note that $D = 75$ is not eligible, since $75 = 3 \cdot 5^2$. Non-eligible numbers in the interval $[3, 103]$ are $\{27, 63, 75, 99\}$.

CFG is interested in values of D for which $h(D)/\sqrt{D}$ is large. Your role is to write a program to help CFG finding these record numbers.

Input

You are given an input file consisting of several test cases, each of them consists of three integers on a single line:

$Dmin \ Dmax \ K$

where $3 \leq Dmin \leq Dmax < 2^{31}$ and are of the form $4k + 3$. Moreover, $Dmax - Dmin \leq 10^6$ and $K < 10^4$. For such values, one has $h(D) < 2^{31}$.

Output

For each test case, your program must determine the eligible values of D in the interval $[Dmin, Dmax]$ for which

$$f(D) = \left\lfloor \frac{1000 h(D)}{\lfloor \sqrt{D} \rfloor} \right\rfloor \geq K.$$

The output will consist of lines:

$D \ h \ f$

where D is a record number, $h = h(D)$ and $f = f(D)$.

If no answer is found, then output a line containing the word 'empty'.

Write a blank line to separate the output of two consecutive cases.

Sample Input

```
3 103 0
27 27 10
```

Sample Output

```
3 1 1000
7 1 500
11 1 333
15 2 666
19 1 250
23 3 750
31 3 600
```

35 2 400
39 4 666
43 1 166
47 5 833
51 2 285
55 4 571
59 3 428
67 1 125
71 7 875
79 5 625
83 3 333
87 6 666
91 2 222
95 8 888
103 5 500

empty